

ON TOTAL FROBENIUS-SCHUR INDICATORS

GONGXIANG LIU AND SIU-HUNG NG

ABSTRACT. We define total Frobenius-Schur indicator for each object in a spherical fusion category $\mathcal{C}$ as a certain canonical sum of its higher indicators. The total indicators are invariants of spherical fusion categories. If $\mathcal{C}$ is the representation category of a semisimple quasi-Hopf algebra H , we prove that the total indicators are non-negative integers which satisfy a certain divisibility condition. In addition, if H is a Hopf algebra, then all the total indicators are positive. Consequently, the positivity of total indicators is a necessary condition for a quasi-Hopf algebra being gauge equivalent to a Hopf algebra. Certain twisted quantum doubles of finite groups and some examples of Tambara-Yamagami categories are discussed for the sufficiency of this positivity condition.

1. INTRODUCTION

The representation category $\text{Rep}(H)$ of a Hopf algebra H is certainly important to the understanding of the algebraic structure of H . The monoidal structure of $\text{Rep}(H)$ has also been playing important roles in other areas of mathematics and physics. For instance, the quantum invariants of knots, links or 3-manifolds constructed from certain Hopf algebras are actually determined by the monoidal structures of their representation categories [33].

Quasi-Hopf algebras are generalizations of Hopf algebras whose representation categories are also monoidal categories. Two quasi-Hopf algebras are said to be *gauge equivalent* if their representation categories are monoidally equivalent. For any finite-dimensional quasi-Hopf algebra K over $\mathbb{C}$, one can obtain another quasi-Hopf algebra K^F by twisting K with a gauge transformation $F \in K \otimes K$ [14], but K^F and K are gauge equivalent. In general, two finite-dimensional quasi-Hopf algebras K and H are gauge equivalent if, and only if, there exists a gauge transformation $F \in K \otimes K$ such that K^F and H are isomorphic as quasi-bialgebras (cf. [7, 25]). However, it is generally difficult to decide the gauge equivalence of two finite-dimensional quasi-Hopf algebras if their Grothendieck rings happen to be isomorphic.

For any finite group G , and a normalized 3-cocycle ω on G with values in $\mathbb{C}^\times$, one can construct a semisimple quasi-Hopf algebra $D^\omega(G)$, called a *twisted quantum double* of G [5]. Dijkgraaf, Pasquier and Roche have asked the question whether $D^\omega(G)$ is gauge equivalent to some Hopf algebra in [5]. One can even ask a more general question: How can one determine whether a given finite-dimensional quasi-Hopf algebra K over $\mathbb{C}$ is gauge equivalent to some Hopf algebra? In view of the reconstruction theorem, the question is equivalent to ask for the existence of a

The first author was partially supported by the NSF grant number 11071111 and the second author was partially supported by NSF grant number DMS1001566.

fibration, i.e. a $\mathbb{C}$ -linear, faithful and exact monoidal functor $\mathcal{F} : \text{Rep}(K) \rightarrow \text{Vect}_{\mathbb{C}}$, where $\text{Vect}_{\mathbb{C}}$ denotes the category of finite-dimensional $\mathbb{C}$ -linear spaces.

This simply stated question is generally difficult to answer. One can work out the simplest example $D^{\omega}(\mathbb{Z}_2)$, where ω is a non-trivial 3-cocycle of $\mathbb{Z}_2$. It is not completely obvious that $D^{\omega}(\mathbb{Z}_2)$ is not gauge equivalent to any Hopf algebra [23].

The n -th Frobenius-Schur (FS) indicator $\nu_n(V)$ of a representation V of a finite group was introduced for more than a century. It has been generalized to the representations of a semisimple (quasi-)Hopf algebra [16, 24], to the primary fields of a rational conformal field theory [3], and more generally to the objects of a pivotal categories [27]. These indicators are preserved by equivalences of pivotal category [loc. cit.]. The arithmetical properties of the FS indicators also encrypt the structure of the underlying tensor categories as well as the quasi-Hopf algebras. For instance, the indicators of any complex representation of a finite group are integers.

If V is an object of a spherical fusion category $\mathcal{C}$ over $\mathbb{C}$, it has been shown that the sequence $\nu(V) := \{\nu_n(V)\}_{n \in \mathbb{N}}$ of FS indicators is a periodic sequence of cyclotomic integers. Moreover, there exists a global period N for all the higher indicator sequences [26]. This global period, denoted by $\text{FSexp}(\mathcal{C})$, is called the Frobenius-Schur exponent of $\mathcal{C}$, and it is also an invariant of spherical fusion categories. The representation category $\text{Rep}(H)$ of a semisimple quasi-Hopf algebra over $\mathbb{C}$ is a spherical fusion category [8], and we simply denote the Frobenius-Schur exponent of $\text{Rep}(H)$ by $\text{FSexp}(H)$.

The FS indicators of a representation of semisimple Hopf algebra are not necessarily integers (see [13] for example). However, if we consider the total indicator $\bar{\nu}(V)$ defined for any object V in a spherical fusion category $\mathcal{C}$ by

$$\bar{\nu}(V) := \sum_{n=1}^N \nu_n(V)$$

where $N = \text{FSexp}(\mathcal{C})$, then we have the following integrality and divisibility theorem for quasi-Hopf algebras.

Theorem A. *Let H be a semisimple quasi-Hopf algebra over $\mathbb{C}$. For any finite-dimensional H -module V , $\bar{\nu}(V)$ is a non-negative integer which satisfies the divisibility*

$$\text{FSexp}(H) \mid (\dim H) \cdot \bar{\nu}(V).$$

In addition, for semisimple Hopf algebras, we have obtained the positivity of total indicators:

Theorem B. *Let H be a semisimple Hopf algebra $\mathbb{C}$. Then, $\bar{\nu}(V) \geq \frac{N \dim V}{\dim H}$ for all finite-dimensional H -modules V , where $N = \text{FSexp}(H)$. In particular, $\bar{\nu}(V) > 0$ for any non-zero H -module V .*

Since $\text{FSexp}(H) = \exp(H)$ when H is a semisimple Hopf algebra, Theorem A provides another perspective for a conjecture of Kashina [11, 12]: $\exp(H)$ divides $\dim H$ for any semisimple Hopf algebra H over $\mathbb{C}$. Moreover, Theorem B yields a necessary condition for a semisimple quasi-Hopf algebra being gauge equivalent to a Hopf algebra. However, this necessary condition may not be sufficient in general. There are integral Tambara-Yamagami categories which has no fibration but their total indicators are all positive. Nevertheless, the positivity of total indicators is a

necessary and sufficient for abelian twisted quantum doubles being gauge equivalent to Hopf algebras.

The organization of the paper is as follows: In Section 2, we introduce the definition of total Frobenius-Schur indicators and prove Theorem A. Section 3 is devoted to the proof of Theorem B. In Section 4, we consider twisted quantum doubles of $D^\omega(G)$ of a finite abelian group G , and show that if $D^\omega(G)$ is commutative, then $D^\omega(G)$ is gauge equivalent to a Hopf algebra if, and only if, $\bar{\nu}(V) > 0$ for all $V \in \text{Rep}(D^\omega(G))$. This provides an answer to a question of Dijkgraaf, Pasquier and Roche [5, p69]. In Section 5, we compute the total indicators for some integral Tambara-Yamagami fusion categories, and we characterize those admitting a fibration in terms of total indicators. As a consequence, semisimple quasi-Hopf algebras with positive total indicators but not gauge equivalent to any Hopf algebra are found.

2. TOTAL FROBENIUS-SCHUR INDICATORS

Throughout this paper, unless stated otherwise, we will work over the field $\mathbb{C}$ of complex numbers; every monoidal category $\mathcal{C}$ in this paper is assumed to be $\mathbb{C}$ -linear abelian with finite-dimensional Hom-spaces over $\mathbb{C}$ and a *strict* simple unit object $\mathbf{1}_{\mathcal{C}}$. All (quasi-)Hopf algebras are assumed to be semisimple and finite-dimensional over $\mathbb{C}$. We denote by $\text{Rep}(H)$ the $\mathbb{C}$ -linear monoidal category of finite-dimensional representations of a quasi-Hopf algebra H . The unit object of $\text{Rep}(H)$, simply denoted by $\mathbf{1}_H$, is the H -module $\mathbb{C}$ equipped with the trivial H -action.

In this section, we collect some conventions, and recall some basic definitions and facts for the discussions in the remaining sections. The readers may refer to [2, 8, 21] for the basic theory of tensor categories and [16, 13, 27, 25] for Frobenius-Schur indicators. The aim of this section is to introduce the definition of total Frobenius-Schur indicators (abbr. total indicators), and to prove Theorem A.

Let $\mathcal{C}$ be a left rigid monoidal category with tensor product $\otimes$. The left dual of $V \in \mathcal{C}$ is a triple $(V^*, \text{db}, \text{ev})$ in which $V^* \in \mathcal{C}$, and $\text{db} : \mathbf{1} \rightarrow V \otimes V^*$ and $\text{ev} : V^* \otimes V \rightarrow \mathbf{1}$ are respectively the associated dual basis and evaluation morphisms of the left dual. The left duality on $\mathcal{C}$ can be extended to a monoidal equivalence $(-)^* : \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$ and hence $(-)^{**} : \mathcal{C} \rightarrow \mathcal{C}$ defines a monoidal equivalence. A pivotal structure on $\mathcal{C}$ is an isomorphism $j : \text{Id} \rightarrow (-)^{**}$ of monoidal functors, and the pair $(\mathcal{C}, j)$ is called a *pivotal category*. We will simply say that $\mathcal{C}$ is a pivotal category when the pivotal structure is understood without ambiguity.

In a pivotal category $(\mathcal{C}, j)$, one can define the left and right pivotal traces for any endomorphism $f : V \rightarrow V$ in $\mathcal{C}$ as

$$\begin{aligned} \underline{\text{ptr}}^r(f) &:= \left(\mathbf{1} \xrightarrow{\text{db}} V \otimes V^* \xrightarrow{f \otimes V^*} V \otimes V^* \xrightarrow{j_V \otimes V^*} V^{**} \otimes V^* \xrightarrow{\text{ev}} \mathbf{1} \right), \\ \underline{\text{ptr}}^\ell(f) &:= \left(\mathbf{1} \xrightarrow{\text{db}} V^* \otimes V^{**} \xrightarrow{V^* \otimes j_V^{-1}} V^* \otimes V \xrightarrow{V^* \otimes f} V^* \otimes V \xrightarrow{\text{ev}} \mathbf{1} \right) \end{aligned}$$

respectively. Note that these traces are scalars as $\mathbf{1}$ is simple. A *spherical category* is a pivotal category $\mathcal{C}$ in which $\underline{\text{ptr}}^r(f) = \underline{\text{ptr}}^\ell(f)$ for all endomorphisms $f \in \mathcal{C}$. In this case, we simply write $\underline{\text{ptr}}$ for the functions $\underline{\text{ptr}}^r$ as well as $\underline{\text{ptr}}^\ell$, and $d(V) := \underline{\text{ptr}}(\text{id}_V)$ is called the *pivotal dimension* of V . In addition, if $\mathcal{C}$ is semisimple with finitely many simple objects up to isomorphism, then $\mathcal{C}$ is called a *spherical fusion category* (cf. [8] for more details on fusion categories). In this case, $d(V)$ is a non-zero real

number (cf. [8]), and the *global dimension* $\dim \mathcal{C}$ of $\mathcal{C}$ is defined as

$$\dim \mathcal{C} = \sum_{V \in \text{Irr}(\mathcal{C})} d(V)^2$$

where $\text{Irr}(\mathcal{C})$ denotes a complete set of non-isomorphic simple objects of $\mathcal{C}$.

By Müger [22], the center $\mathcal{Z}(\mathcal{C})$ is a *modular tensor category*. In particular, the associated *twist* (or *ribbon structure*) θ has finite order [2, 34]. Moreover, the forgetful functor $F : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$ admits a two-sided adjoint $K : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C})$.

Let V be an object in a pivotal category $(\mathcal{C}, j)$ with associativity isomorphism Φ . We write $V^{\otimes n}$ for the n -fold tensor power of $V \in \mathcal{C}$ with rightmost parentheses. By coherence theorem, there is a unique isomorphism

$$\Phi^{(n)} : V^{\otimes(n-1)} \otimes V \rightarrow V^{\otimes n}$$

which is a composition of tensor products of id , Φ and Φ^{-1} . One can define the $\mathbb{C}$ -linear operator $E_V^{(n)} : \mathcal{C}(\mathbf{1}, V^{\otimes n}) \rightarrow \mathcal{C}(\mathbf{1}, V^{\otimes n})$ by setting

$$\begin{aligned} E_V^{(n)}(f) := & \left(\mathbf{1} \xrightarrow{\text{db}} V^* \otimes V^{**} \xrightarrow{(V^* \otimes f) \otimes V^{**}} (V^* \otimes V^{\otimes n}) \otimes V^{**} \xrightarrow{\Phi^{-1} \otimes j^{-1}} \right. \\ & \left. ((V^* \otimes V) \otimes V^{\otimes(n-1)}) \otimes V \xrightarrow{(\text{ev} \otimes V^{\otimes(n-1)}) \otimes V} V^{\otimes(n-1)} \otimes V \xrightarrow{\Phi^{(n)}} V^{\otimes n} \right). \end{aligned}$$

Following [27, Sect. 3], the n -th *Frobenius-Schur indicator* $\nu_n(V)$ of V is defined as the scalar

$$\nu_n(V) = \text{Tr}(E_V^{(n)}).$$

These indicators are proved to be invariant under pivotal equivalences, and $\nu_n(V)$ is a cyclotomic integer in $\mathbb{Q}(\zeta_n)$ where ζ_n is a primitive n -th root of unity (cf. [27]).

Since the antipode of a semisimple Hopf algebra H is an involution [15], the representation category $\text{Rep}(H)$ is a spherical fusion category in which the pivotal structure is the usual canonical isomorphism $j_V : V \rightarrow V^{**}$ of finite-dimensional vector spaces. In this case, the pivotal dimension $d(V)$ of $V \in \text{Rep}(H)$ is the ordinary dimension of V . More generally, for a semisimple quasi-Hopf algebra H , there is a unique (spherical) pivotal structure on $\text{Rep}(H)$ such that $d(V)$ is the ordinary dimension of V for all $V \in \text{Rep}(H)$ [8]. Moreover, $\nu_n(V)$ can be expressed in terms of the associator, the quasi-antipode and the normalized integral Λ of H (cf. [25, Sect. 4]). When H is a Hopf algebra, we recover the n -th Frobenius-Schur indicator formula of V introduced in [16]:

$$(2.1) \quad \nu_n(V) = \chi_V(\Lambda^{[n]})$$

where χ_V is the character of H afforded by V , Λ is the normalized integral of H and $\Lambda^{[n]} = \Lambda_1 \Lambda_2 \cdots \Lambda_n$. Here, $x_1 \otimes \cdots \otimes x_n$ denotes the Sweedler notation of the n -fold comultiplication of $x \in H$.

The *Frobenius-Schur exponent*, denoted by $\text{FSexp}(\mathcal{C})$, of a spherical category $\mathcal{C}$ (cf. [26]) is the least positive integer n such that $\nu_n(V) = d(V)$ for all $V \in \mathcal{C}$. If such an integer does not exist, $\text{FSexp}(\mathcal{C})$ is defined to be ∞ . However, the Frobenius-Schur exponent of a spherical fusion category is always finite because the following theorem proved in [26, Thm. 4.1 and 5.5].

Theorem 2.1. *Let $\mathcal{C}$ be a spherical fusion category, θ the twist of $\mathcal{Z}(\mathcal{C})$ and $K : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C})$ the two-sided adjoint of the forgetful functor $F : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$. Then, for $V \in \mathcal{C}$,*

- (i) $\nu_n(V) = \frac{1}{\dim \mathcal{C}} \underline{\text{ptr}}(\theta_{K(V)})$ for all $V \in \mathcal{C}$, and
- (ii) $\text{FSexp}(\mathcal{C}) = \text{ord}(\theta)$. □

For a simple object $V \in \mathcal{C}$, it is known that $\nu_1(V) = 1$ if $V \cong \mathbf{1}$, and 0 otherwise. The second indicator of V can only be 0, 1, -1 depending on whether V is self-dual or not. By Theorem 2.1, we know $\nu_N(V) = d(V)$ if $N = \text{FSexp}(\mathcal{C})$. The meaning of higher indicators of V are more obscure and they are not rational integers in general (cf. [13, Ex. 7.5]). The theorem implies that the indicator sequence $\nu(V) := \{\nu_n(V)\}_{n \in \mathbb{N}}$ of V is periodic for any object V of a spherical fusion category $\mathcal{C}$. Moreover, $\text{FSexp}(\mathcal{C})$ is the global period of all the indicator sequences of $\mathcal{C}$. The average value or the sum of these indicators over a period should also be an important invariant.

Definition 2.2. Let $\mathcal{C}$ be a spherical fusion category and $N = \text{FSexp}(\mathcal{C})$. The total Frobenius-Schur indicator of $V \in \mathcal{C}$, denoted by $\bar{\nu}(V)$, is defined as

$$\bar{\nu}(V) := \sum_{i=1}^N \nu_i(V).$$

To prove Theorem A, we first derive a formula for the total indicator $\bar{\nu}(V)$ of an object V in a spherical fusion category $\mathcal{C}$ in terms of some data of the center $\mathcal{Z}(\mathcal{C})$. Recall from [22, Proposition 8.1] that the forgetful functor $F : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$ has a two-sided adjoint $K : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C})$. It follows from the semisimplicity of $\mathcal{C}$, we have

$$K(V) \cong \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} \dim(\mathcal{Z}(\mathcal{C})(K(V), X)) X$$

for $V \in \mathcal{C}$, where $\text{Irr}(\mathcal{Z}(\mathcal{C}))$ is a complete set of non-isomorphic simple objects of $\mathcal{Z}(\mathcal{C})$. For simplicity, we set

$$[U : V]_{\mathcal{C}} := \dim(\mathcal{C}(U, V)) \quad \text{for all } U, V \in \mathcal{C}.$$

Since K is a left adjoint of F , $\mathcal{Z}(\mathcal{C})(K(V), X) \cong \mathcal{C}(V, F(X))$ and so

$$[K(V) : X]_{\mathcal{Z}(\mathcal{C})} = [V : F(X)]_{\mathcal{C}}$$

for $X \in \mathcal{Z}(\mathcal{C})$ and $V \in \mathcal{C}$.

Since $\dim(\mathcal{C}(U, V)) = \dim(\mathcal{C}(V, U)) = \dim(\mathcal{C}(U^*, V^*))$, we have

$$[V : U]_{\mathcal{C}} = [U : V]_{\mathcal{C}} = [U^*, V^*]_{\mathcal{C}} \text{ for all } U, V \in \mathcal{C}.$$

Thus, we have

$$(2.2) \quad K(V) \cong \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X) : V]_{\mathcal{C}} X.$$

Let θ be the twist of $\mathcal{Z}(\mathcal{C})$. For any $X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))$, we define $\omega_X \in \mathbb{C}$ by the equation

$$\theta_X = \omega_X \text{id}_X.$$

Note that ω_X is an N -th root of unity by Theorem 2.1.

Proposition 2.3. Let $\mathcal{C}$ be a spherical fusion categories over $\mathbb{C}$ with $\text{FSexp}(\mathcal{C}) = N$, and $V \in \mathcal{C}$. Then

- (i) $\bar{\nu}(V)$ is an algebraic integer invariant under pivotal equivalence, i.e. if $\mathcal{F} : \mathcal{C} \rightarrow \mathcal{D}$ defines an equivalence of pivotal categories, then $\bar{\nu}(V) = \bar{\nu}(\mathcal{F}(V))$.

(ii) Moreover,

$$(2.3) \quad \bar{\nu}(V) = \frac{N}{\dim(\mathcal{C})} \sum_{\substack{X \in \text{Irr}(\mathcal{Z}(\mathcal{C})) \\ \theta_X = \text{id}_X}} [F(X) : V]_{\mathcal{C}} d(X)$$

where $F : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$ is the forgetful functor.

Proof. Statement (i) is an immediate consequence of the fact that $\nu_n(V)$ are algebraic integers for all n , and that both $\nu_n(V)$ and $\text{FSexp}(\mathcal{C})$ are invariant under pivotal equivalences.

(ii) By Theorem 2.1, we have

$$\nu_n(V) = \frac{1}{\dim(\mathcal{C})} \text{ptr}(\theta_K^n(V)) = \frac{1}{\dim(\mathcal{C})} \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X) : V]_{\mathcal{C}} \omega_X^n d(X).$$

Therefore,

$$\begin{aligned} \bar{\nu}(V) &= \sum_{i=1}^N \nu_i(V) \\ &= \frac{1}{\dim(\mathcal{C})} \sum_{i=1}^N \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X) : V]_{\mathcal{C}} \omega_X^i d(X) \\ &= \frac{1}{\dim(\mathcal{C})} \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X) : V]_{\mathcal{C}} \sum_{i=1}^N \omega_X^i d(X) \\ &= \frac{N}{\dim(\mathcal{C})} \sum_{\substack{X \in \text{Irr}(\mathcal{Z}(\mathcal{C})) \\ \omega_X = 1}} [F(X) : V]_{\mathcal{C}} d(X). \end{aligned}$$

Here, the last equality follows from the fact that ω_X is an N -th root of unity. $\square$

We can now prove Theorem A.

Proof of Theorem A. Let H be a semisimple quasi-Hopf algebra. Consider the canonical pivotal structure on $\text{Rep}(H)$. Then $\text{Rep}(H)$ is a spherical fusion category with $d(V) = \dim V$ for all $V \in \text{Rep}(H)$. In particular, $\dim(\text{Rep}(H)) = \dim H$ and $d(X)$ is a non-negative integer for all $X \in \mathcal{Z}(\text{Rep}(H))$. It follows from Proposition 2.3 (ii) that

$$\bar{\nu}(V) = \frac{N}{\dim H} \sum_{\substack{X \in \text{Irr}(\mathcal{Z}(\mathcal{C})) \\ \omega_X = 1}} [F(X) : V]_{\mathcal{C}} d(X)$$

is a non-negative rational number. By Proposition 2.3 (i), $\bar{\nu}(V)$ is a non-negative integer. Since

$$\frac{\bar{\nu}(V) \dim H}{N} = \sum_{\substack{X \in \text{Irr}(\mathcal{Z}(\mathcal{C})) \\ \omega_X = 1}} [F(X) : V]_{\mathcal{C}} d(X) \in \mathbb{Z},$$

we establish the divisibility $N \mid (\dim H) \bar{\nu}(V)$. $\square$

If H is a semisimple Hopf algebra, $\exp(H) = \text{ord}(\theta) = \text{FSexp}(H)$ (cf. [6, Thm 2.5] and Theorem 2.1). Therefore, Theorem A is related to the following well-known conjecture considered by Kashina [11, 12].

Conjecture 2.4. *Let H be a semisimple Hopf algebra over $\mathbb{C}$. Then the exponent of H divides $\dim(H)$.*

By the Cauchy theorem for Hopf algebras [13, Sect. 6] (see so [26, Thm. 8.4]), $\dim H$ and $\exp(H)$ have the same prime factors. Thus, if $\gcd\{\bar{\nu}(V) \mid V \in \text{Irr}(H)\}$ is relatively prime to $\dim H$, then the conjecture will be proved for H . However, the Kac algebra K of dimension 8 is an example in which $\gcd\{\bar{\nu}(V) \mid V \in \text{Irr}(K)\} = 4$ (see Example 3.6). Nevertheless, any upper bound of $\gcd\{\bar{\nu}(V) \mid V \in \text{Irr}(H)\}$ will shed light on the conjecture.

By Theorem A, $\bar{\nu}(V) = 0$ is an extreme value and it is possible for some quasi-Hopf algebra demonstrated in the following example.

Example 2.5. Let G be a finite group and ω a normalized 3-cocycle on G with coefficients in $\mathbb{C}^\times$. Following [25, Sect. 7], one can construct a quasi-Hopf algebra $H(G, \omega) = (\mathbb{C}[G]^*, \Delta, \varepsilon, \phi, \alpha, \beta, S)$ where multiplication, identity, comultiplication Δ , counit ε , and antipode S are the same as the structure maps of the dual $\mathbb{C}[G]^*$ of the group algebra $\mathbb{C}[G]$, and ϕ, α , and β are given by

$$\phi = \sum_{a,b,c \in G} \omega(a,b,c) e(a) \otimes e(b) \otimes e(c), \quad \alpha = 1, \quad \text{and} \quad \beta = \sum_{a \in G} \omega(a, a^{-1}, a)^{-1} e(a),$$

where $\{e(x) \mid x \in G\}$ is the dual basis of G for $\mathbb{C}[G]^*$. If $\omega' \in Z^3(G, \mathbb{C}^\times)$ is cohomologous to ω , then $H(G, \omega)$ can be twisted to the quasi-bialgebra $H(G, \omega')$ by a gauge transformation (see [14, XV] for gauge equivalence of quasi-bialgebras). In particular, if ω is a coboundary of the 2-cochain $f : G \times G \rightarrow \mathbb{C}^\times$, then $H(G, \omega)$ can be twisted by the gauge transformation $F = \sum_{a,b \in G} f(a,b) e(a) \otimes e(b)$ to the ordinary bialgebra $\mathbb{C}[G]^*$.

We now consider a special case when G is of order 2. Let $G = \{1, x\}$ be an abelian group of order 2 and ω a 3-cocycle of G given by

$$\omega(a,b,c) = \begin{cases} -1 & \text{if } a = b = c = x, \\ 1 & \text{otherwise.} \end{cases}$$

Then $H = H(G, \omega)$ is a 2-dimensional commutative quasi-Hopf algebra. Let V be the nontrivial simple H -module. As computed in [26, Ex. 5.4], $\text{FSexp}(H) = 4$ and $\nu_n(V) = \cos(\frac{n\pi}{2})$. Therefore, $\bar{\nu}(V) = 0$.

3. PROOF OF THEOREM B

Example 2.5 shows that $\bar{\nu}(V) = 0$ for some simple module V of a semisimple quasi-Hopf algebra. However, this cannot happen for semisimple Hopf algebras which is stated in Theorem B. We will prove this theorem in this section.

Recall that the Drinfeld double $D(H)$ of a finite-dimensional Hopf algebra H is the bicrossproduct $H^{*\text{cop}} \bowtie H$ (cf. [19, 14]). In particular, $D(H)$ is a finite-dimensional Hopf algebra with comultiplication and multiplication of $D(H)$ described by

$$\Delta(f \bowtie a) = (f_2 \bowtie a_1) \otimes (f_1 \bowtie a_2)$$

and

$$(f \bowtie a)(g \bowtie b) = f(g(S^{-1}(a_3)?a_1)) \bowtie a_2b$$

respectively, for $f, g \in H^*$ and $a, b \in H$. Here $a_1 \otimes a_2$ and $f_1 \otimes f_2$ are respectively the Sweedler notation for the comultiplications of $a \in H$ and $f \in H^*$ with the

summation suppressed. The Hopf algebra $D(H)$ admits a canonical universal $\mathcal{R}$ -matrix $R = \sum_i \varepsilon \bowtie h_i \otimes h^i \bowtie 1$ where $\{h_i\}_i$ and $\{h^i\}_i$ are dual bases of H and H^* respectively.

It should be well-known to experts that $\text{Rep}(D(H))$ and $\text{Rep}(D(H^*))$ are equivalent braided monoidal categories. For the sake of completeness, we provide a proof of the statement in terms of gauge equivalence for subsequent discussion.

Lemma 3.1. *Let H be a finite-dimensional Hopf algebra over any field $\mathbb{k}$, not necessarily semisimple. Then $D(H)$ and $D(H^*)$ are gauge equivalent quasi-triangular Hopf algebras via the gauge transformation $R \in D(H^*) \otimes D(H^*)$, which is the universal $\mathcal{R}$ -matrix of $D(H^*)$, and the algebra isomorphism*

$$\sigma : D(H) \rightarrow D(H^*), \quad f \bowtie a \mapsto (1 \bowtie f)(a \bowtie \varepsilon).$$

Here, H^{**} is identified with H as Hopf algebras under the natural isomorphism $j : H \rightarrow H^{**}$ of vector spaces.

Proof. Since $\Delta_{D(H^*)}^{\text{op}}(x) = R\Delta_{D(H^*)}(x)R^{-1}$ for all $x \in D(H^*)$, it suffices to show that $\sigma : D(H) \rightarrow D(H^*)^{\text{cop}}$ is a bialgebra homomorphism, but this is straightforward verification. $\square$

The algebra isomorphism $\sigma : D(H) \rightarrow D(H^*)$ defines a braided monoidal equivalence $(\mathcal{F}_\sigma, \xi, \text{id}) : \text{Rep}(D(H^*)) \rightarrow \text{Rep}(D(H))$ (cf. [14, XIII.3.2]). Here, $\mathcal{F}_\sigma : \text{Rep}(D(H^*)) \rightarrow \text{Rep}(D(H))$ is the $\mathbb{C}$ -linear functor with $\mathcal{F}_\sigma(V)$ defined as the $D(H)$ -module via the isomorphism σ for $V \in \text{Rep}(D(H^*))$, and $\mathcal{F}_\sigma : \text{Hom}_{D(H^*)}(V, W) \rightarrow \text{Hom}_{D(H)}(\mathcal{F}_\sigma V, \mathcal{F}_\sigma W)$ the identity function. The coherence isomorphism $\xi : \mathcal{F}_\sigma(V) \otimes \mathcal{F}_\sigma(W) \rightarrow \mathcal{F}_\sigma(V \otimes W)$ is the left action of R^{-1} , and $\mathcal{F}_\sigma(\mathbf{1}_{D(H^*)}) = \mathbf{1}_{D(H)}$.

The Hopf algebra H^* can be considered as a subalgebra of $D(H)$ and $D(H^*)$ via the embeddings $i_1 : H^* \rightarrow D(H^*)$ and $i_2 : H^* \rightarrow D(H)$ defined by

$$i_1(f) = 1 \bowtie f, \quad i_2(f) = f \bowtie 1$$

respectively, for $f \in H^*$. It follows immediately from the definition of σ that we have the commutative diagram of algebra maps:

$$(3.1) \quad \begin{array}{ccc} D(H) & \xrightarrow{\sigma} & D(H^*) \\ i_2 \uparrow & \nearrow i_1 & \\ H^* & & \end{array}$$

This implies the following lemma for the two pairs of induction and restriction functors.

Lemma 3.2. *Let $\text{Res}_{H^*}^{D(H^*)}$ and $\text{Res}_{H^*}^{D(H)}$ be the restriction functors along the embeddings i_1 and i_2 respectively, and $\text{Ind}_{H^*}^{D(H^*)}$ and $\text{Ind}_{H^*}^{D(H)}$ the associated induction functors. Then we have*

$$(3.2) \quad \mathcal{F}_\sigma \circ \text{Ind}_{H^*}^{D(H^*)} = \text{Ind}_{H^*}^{D(H)} \quad \text{and} \quad \text{Res}_{H^*}^{D(H)} \circ \mathcal{F}_\sigma = \text{Res}_{H^*}^{D(H^*)}. \quad \square$$

The preceding lemmas hold for any finite-dimensional Hopf algebras. We now turn to semisimple Hopf algebras H . In this case, $\mathcal{C} = \text{Rep}(H)$ is a spherical fusion category, and the *right* center $\mathcal{Z}(\mathcal{C})$ of $\mathcal{C}$ is equivalent to the $\text{Rep}(D(H))$ as braided monoidal category. Equipped with the canonical pivotal structure, $\text{Rep}(D(H))$ is a modular tensor category with the twist θ given by the action of the Drinfeld

element of $D(H)$. The forgetful functor $F : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$ is the restriction $\text{Res}_H^{D(H)}$ which has a left adjoint $\text{Ind}_H^{D(H)} : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C})$. Thus, by the uniqueness of adjoint functor, $K : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$ is equivalent to $\text{Ind}_H^{D(H)}$. If V is a self-dual H -module, then so is $K(V)$. This observation even holds for spherical fusion categories.

Lemma 3.3. *Let $\mathcal{C}$ be a spherical fusion category, $\mathcal{Z}(\mathcal{C})$ the center of $\mathcal{C}$ and $K : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C})$ the left adjoint of the forgetful functor $F : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$. Then $K(V)$ is a self-dual object of $\mathcal{Z}(\mathcal{C})$ whenever $V \in \mathcal{C}$ is self-dual. In particular, $K(\mathbf{1}_{\mathcal{C}})$ is self-dual. Moreover, if θ is the twist of $\mathcal{Z}(\mathcal{C})$, then $\theta_{K(\mathbf{1}_{\mathcal{C}})} = \text{id}_{K(\mathbf{1}_{\mathcal{C}})}$.*

Proof. Note that for $X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))$, X^* is isomorphic to a unique object of $\text{Irr}(\mathcal{Z}(\mathcal{C}))$. Let $V \in \mathcal{C}$ be self-dual. It follows from (2.2) that

$$\begin{aligned} K(V)^* &\cong \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X) : V]_{\mathcal{C}} X^* \\ &\cong \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X^*) : V]_{\mathcal{C}} X \\ &\cong \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X)^* : V]_{\mathcal{C}} X \\ &\cong \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X)^* : V^*]_{\mathcal{C}} X \\ &\cong \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X) : V]_{\mathcal{C}} X \\ &\cong K(V). \end{aligned}$$

Here, the third isomorphism is a consequence of the fact that the forgetful functor $F : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$ defines a monoidal functor, and the last isomorphism follows from the remark preceding Proposition 2.3. Since $\mathbf{1}_{\mathcal{C}}$ is self-dual, and so is $K(\mathbf{1}_{\mathcal{C}})$.

By [28, Prop. 2.8 (iii)], $[F(X) : \mathbf{1}_{\mathcal{C}}]_{\mathcal{C}} \neq 0$ implies $\omega_X = 1$. Therefore,

$$\theta_{K(\mathbf{1}_{\mathcal{C}})} = \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X) : \mathbf{1}_{\mathcal{C}}]_{\mathcal{C}} \theta_X = \sum_{X \in \text{Irr}(\mathcal{Z}(\mathcal{C}))} [F(X) : \mathbf{1}_{\mathcal{C}}]_{\mathcal{C}} \text{id}_X = \text{id}_{K(\mathbf{1}_{\mathcal{C}})}.$$

This proves the last assertion. $\square$

We can now prove Theorem B.

Proof of Theorem B. Let $\mathcal{C} = \text{Rep}(H)$. In view of Proposition 2.3 (ii), it suffices to show the inequality

$$\sum_{\substack{X \in \text{Irr}(D(H)) \\ \theta_X = \text{id}_X}} [\text{Res}_H^{D(H)} X : V]_{\mathcal{C}} \dim(X) \geq \dim(V)$$

for all $V \in \mathcal{C}$. This is equivalent to show that $[\text{Res}_H^{D(H)} W : V]_{\mathcal{C}} > 0$ for some $D(H)$ -module W satisfying $\theta_W = \text{id}_W$.

Recall that H^* is also a semisimple Hopf algebra [15]. By Lemma 3.3, $Y = \text{Ind}_{H^*}^{D(H^*)}(\mathbf{1}_{H^*})$ is a self-dual $D(H^*)$ -module and $\theta_Y = \text{id}_Y$. Moreover,

$$(3.3) \quad Y \cong \sum_{U \in \text{Irr}(D(H^*))} [U : \mathbf{1}_{H^*}]_{\mathcal{C}} U$$

where $\mathcal{C}' = \text{Rep}(H^*)$. Since $\text{Rep}(D(H^*))$ and $\text{Rep}(D(H))$ are spherical categories equipped with their canonical pivotal structures, $\mathcal{F}_\sigma : \text{Rep}(D(H^*)) \rightarrow \text{Rep}(D(H))$ defines an equivalence of braided monoidal categories as well as pivotal categories. Therefore, by [9, Prop. 6.2], we have

$$\text{id}_{\mathcal{F}_\sigma Y} = \mathcal{F}_\sigma(\text{id}_Y) = \mathcal{F}_\sigma(\theta_Y) = \theta_{\mathcal{F}_\sigma Y}.$$

In particular, $W = \mathcal{F}_\sigma Y$ is a self-dual $D(H)$ -module satisfying $\theta_W = \text{id}_W$. Since $\text{Ind}_H^{D(H)}$ is a left adjoint of $\text{Res}_H^{D(H)}$, we have

$$[V : \text{Res}_H^{D(H)} W]_{\mathcal{C}} = [\text{Ind}_H^{D(H)} V : W]_{\mathcal{Z}(\mathcal{C})}.$$

However, by Lemma 3.2, $W = \text{Ind}_{H^*}^{D(H)} \mathbf{1}_{H^*}$. Therefore,

$$\begin{aligned} \text{Hom}_{D(H)} \left(\text{Ind}_H^{D(H)} V, W \right) &= \text{Hom}_{D(H)} \left(\text{Ind}_H^{D(H)} V, \text{Ind}_{H^*}^{D(H)} \mathbf{1}_{H^*} \right) \\ &\cong \text{Hom}_{D(H)} \left(\left(\text{Ind}_{H^*}^{D(H)} \mathbf{1}_{H^*} \right)^* \otimes \left(\text{Ind}_H^{D(H)} V \right), \mathbf{1}_{D(H)} \right) \\ &\cong \text{Hom}_{D(H)} \left(\left(\text{Ind}_{H^*}^{D(H)} \mathbf{1}_{H^*} \right) \otimes \left(\text{Ind}_H^{D(H)} V \right), \mathbf{1}_{D(H)} \right). \end{aligned}$$

By [4, Thm. 8 (1)], we have

$$\left(\text{Ind}_{H^*}^{D(H)} \mathbf{1}_{H^*} \right) \otimes \left(\text{Ind}_H^{D(H)} V \right) \cong D(H)^{\dim V}$$

as $D(H)$ -modules. Thus,

$$[\text{Ind}_H^{D(H)} V : W]_{\mathcal{Z}(\mathcal{C})} = [D(H)^{\dim V} : \mathbf{1}_{D(H)}]_{\mathcal{Z}(\mathcal{C})} = \dim V,$$

and so $[V : \text{Res}_H^{D(H)} W]_{\mathcal{C}} = \dim V$. This completes the proof. $\square$

Definition 3.4. A quasi-Hopf algebra H is said to be *genuine* if H is not gauge equivalent to any ordinary Hopf algebra.

Theorem B provides a sufficient condition for a semisimple quasi-Hopf algebra being genuine.

Corollary 3.5. *Let H be a semisimple quasi-Hopf algebra. If there exists $V \in \text{Rep}(H)$ such that $\bar{\nu}(V) = 0$, then H is a genuine quasi-Hopf algebra.*

Proof. Assume contrary. Then H is gauge equivalent to a Hopf algebra H' . There exists a monoidal equivalence $\mathcal{F} : \text{Rep}(H) \rightarrow \text{Rep}(H')$ (see [25, Thm. 2.2]). Thus, $\mathcal{F}$ preserves their canonical pivotal structures (cf. [27]). By Proposition 2.3 (i) and Theorem B, we have

$$\bar{\nu}(V) = \bar{\nu}(\mathcal{F}(V)) > 0$$

for all $V \in \text{Rep}(H)$. $\square$

The following example shows that the existence of vanishing total indicators can also be a necessary condition for a genuine quasi-Hopf algebra.

Example 3.6. There are exactly four gauge inequivalent 8-dimensional quasi-Hopf algebras with five simple objects $\{a_1, a_2, a_3, a_4, m\}$ and fusion rules given by

$$\{a_1, a_2, a_3, a_4\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2, \quad m^2 \cong \sum_{i=1}^4 a_i \quad \text{and} \quad ma_i \cong a_i m \cong m$$

for $i = 1, 2, 3, 4$. Their categories of representations are Tambara-Yamagami categories [31]. They are $\mathbb{C}[Q]$, $\mathbb{C}[D]$, the 8-dimensional Kac algebra K and its twisted

version K_u (see [25, Sec.6] for details), where Q and D are, respectively, the quaternion group and the dihedral group of order 8. We can assume a_1 to be the unit object. Then $\nu_n(a_1) = 1$ for all $n \geq 1$. The objects a_2, a_3 and a_4 are 1-dimensional representations of these algebras, and their orders are 2. The indicator $\nu_n(a_i) = 1$ if n is even, and 0 otherwise, for $i = 2, 3, 4$. Thus $\bar{\nu}(a_i) > 0$ for $i = 1, 2, 3, 4$. By [25, Sec. 6], we find the following table:

	$\nu_2(m)$	$\nu_3(m)$	$\nu_4(m)$	$\nu_5(m)$	$\nu_6(m)$	$\nu_7(m)$	$\nu_8(m)$	$\bar{\nu}(m)$
K	1	0	0	0	1	0	2	4
K_u	-1	0	0	0	-1	0	2	0
$\mathbb{C}[D]$	1	0	2	0	1	0	2	3
$\mathbb{C}[Q]$	-1	0	2	0	-1	0	2	1

In particular, the Frobenius-Schur exponents of $K, K_u, \mathbb{C}[D], \mathbb{C}[Q]$ are 8, 8, 4, 4 respectively. The quasi-Hopf algebra K_u is the only one in the list for which $\bar{\nu}(m) = 0$. Therefore, K_u is a genuine quasi-Hopf algebra. It is well-known that $K, \mathbb{C}[D], \mathbb{C}[Q]$ are all the noncommutative semisimple Hopf algebras of dimension 8 (cf. [18]). Their corresponding indicators $\nu_n(m)$ can be computed using the formula (2.1).

4. TWISTED QUANTUM DOUBLES

The main result of this section is to show that existence of vanishing total indicators is also a necessary condition for an *abelian* twisted quantum double, considered in [23], being genuine.

The *twisted quantum double* $D^\omega(G)$ of G relative to a normalized 3-cocycle $\omega : G \times G \times G \rightarrow \mathbb{C}^\times$ is the semisimple quasi-Hopf algebra with underlying vector space $\mathbb{C}[G]^* \otimes \mathbb{C}[G]$ in which multiplication, comultiplication Δ , associator ϕ , counit ε and quasi-antipode (S, α, β) are given by

$$\begin{aligned}
(e(g) \otimes x)(e(h) \otimes y) &= \theta_g(x, y) \delta_{g^x, h} e(g) \otimes xy, \\
\Delta(e(g) \otimes x) &= \sum_{hk=g} \gamma_x(h, k) e(h) \otimes x \otimes e(k) \otimes x, \\
\phi &= \sum_{g, h, k \in G} \omega(g, h, k)^{-1} e(g) \otimes 1 \otimes e(h) \otimes 1 \otimes e(k) \otimes 1, \\
\varepsilon(e(g) \otimes x) &= \delta_{g, 1}, \quad \alpha = 1, \quad \beta = \sum_{g \in G} \omega(g, g^{-1}, g) e(g) \otimes 1, \\
S(e(g) \otimes x) &= \theta_{g^{-1}}(x, x^{-1})^{-1} \gamma_x(g, g^{-1})^{-1} e(x^{-1} g^{-1} x) \otimes x^{-1},
\end{aligned}$$

where $\{e(g) | g \in G\}$ is the dual basis of $\{g | g \in G\}$, $\delta_{g, 1}$ is the Kronecker delta, $g^x = x^{-1} g x$, and

$$\begin{aligned}
\theta_g(x, y) &= \frac{\omega(g, x, y) \omega(x, y, (xy)^{-1} g xy)}{\omega(x, x^{-1} g x, y)}, \\
\gamma_g(x, y) &= \frac{\omega(x, y, g) \omega(g, g^{-1} x g, g^{-1} y g)}{\omega(x, g, g^{-1} y g)}
\end{aligned}$$

for any $x, y, g \in G$ (cf. [5, 14]).

As in [23], we denote by Γ^ω the set of all group-like elements of $D^\omega(G)$ and call ω *abelian* if $D^\omega(G)$ is a commutative algebra. This can only happen when G is abelian and Γ^ω spans $D^\omega(G)$. The following theorem characterizes those abelian twisted

doubles of finite groups which are genuine quasi-Hopf algebras, and it provides an answer for a question of Dijkgraaf, Pasquier and Roche posed in [5, p69].

Theorem 4.1. *Let G be a finite abelian group, and ω a normalized 3-cocycle of G such that $D^\omega(G)$ is a commutative algebra. Then $D^\omega(G)$ is a genuine quasi-Hopf algebra if, and only if, there exists $V \in \text{Rep}(D^\omega(G))$ such that $\bar{\nu}(V) = 0$.*

To prove this theorem, we need some preparations. Recall that an Eilenberg-MacLane (EM) 3-cocycle of G is a pair (ω, c) where $\omega \in Z^3(G, \mathbb{C}^\times)$ and c a 2-cochain of G satisfying the following conditions:

$$(4.1) \quad \frac{c(xy, z)}{c(x, z)c(y, z)} \frac{\omega(x, z, y)}{\omega(x, y, z)\omega(z, x, y)} = 1 = \frac{c(x, yz)}{c(x, y)c(x, z)} \frac{\omega(x, y, z)\omega(y, z, x)}{\omega(y, x, z)}$$

for all $x, y, z \in G$. The EM 3-cocycle (ω, c) is a coboundary if there exists a 2-cochain h of G such that

$$\omega = \delta h \quad \text{and} \quad c(x, y) = \frac{h(x, y)}{h(y, x)}.$$

The EM cohomology group $H_{ab}^3(G, \mathbb{C}^\times)$ is then defined by

$$H_{ab}^3(G, \mathbb{C}^\times) = \frac{Z_{ab}^3(G, \mathbb{C}^\times)}{B_{ab}^3(G, \mathbb{C}^\times)}$$

where $Z_{ab}^3(G, \mathbb{C}^\times)$ and $B_{ab}^3(G, \mathbb{C}^\times)$ are respectively the abelian groups of EM 3-cocycles and 3-coboundaries.

For $(\omega, c) \in Z_{ab}^3(G, \mathbb{C}^\times)$, one can assign the function $t(x) := c(x, x)$, called its trace. The trace t of an EM 3-cocycle of G is a quadratic function on G , that means

- (i) $t(x^a) = t(x)^{a^2}$ for $a \in \mathbb{Z}$, and
- (ii) $b_t(x, y) := \frac{t(xy)}{t(x)t(y)}$ defines a bicharacter of G .

The following theorem of Eilenberg and Mac Lane is essential to the discussion in this section (cf. [17, Thm. 3] and [10, §7]).

Theorem 4.2 (Eilenberg-MacLane). *Let $Q(G, \mathbb{C}^\times)$ be the abelian group of all quadratic functions from G to $\mathbb{C}^\times$. The map assigning to each 3-cocycle its trace induces an isomorphism*

$$H_{ab}^3(G, \mathbb{C}^\times) \xrightarrow{\cong} Q(G, \mathbb{C}^\times). \quad \square$$

Remark 4.3. Recall the example of quasi-Hopf algebra $H(G, \omega)$ defined in Example 2.5 for a normalized 3-cocycle ω on a finite group G . Let $\mathcal{C} = \text{Rep}(H(G, \omega))$. Then $\text{Irr}(\mathcal{C})$ consists of a set of 1-dimensional representations V_x indexed by G . Moreover, a pair (ω, c) is an EM 3-cocycle of a finite abelian group G if, and only if, $\text{Rep}(H(G, \omega))$ is a braided spherical fusion category with the braiding given by

$$c_{V_x, V_y} := \left(V_x \otimes V_y \xrightarrow{c(x, y)} V_y \otimes V_x \right)$$

for $x, y \in G$.

For simplicity, we say that a quadratic function $t : G \rightarrow \mathbb{C}^\times$ of a finite abelian group G can be obtained from a bicharacter b of G if $t(x) = b(x, x)$ for all $x \in G$. We denote by $K(G, \mathbb{C}^\times)$ the set of all quadratic functions which can be obtained from some bicharacters of G .

Lemma 4.4. *Let G be a finite abelian group and (ω, c) an EM 3-cocycle of G with trace t . Then:*

- (i) ω is a coboundary if, and only if, $t \in K(G, \mathbb{C}^\times)$.
- (ii) ω^2 is a 3-coboundary of G .

Proof. Let us denote the cohomology class represented by the EM 3-cocycle (ω, c) of G as $[(\omega, c)]$. Then ω is a coboundary if, and only if, $[(\omega, c)] = [(1, b)]$ for some 2-cochain b of G . By (4.1), b is a bicharacter of G and $t(x) = c(x, x) = b(x, x)$ for all $x \in G$. Therefore, $t \in K(G, \mathbb{C}^\times)$. Conversely, suppose there exists a bicharacter b on G such that $b(x, x) = t(x)$ for all $x \in G$. Since $(1, b)$ is an EM 3-cocycle which has the same trace as (ω, c) , $[(\omega, c)] = [(1, b)]$ by Theorem 4.2. In particular, ω is a coboundary. This proves statement (i).

(ii) By (4.1), $b(x, y) := c(x, y)c(y, x)$ defines a bicharacter of G , and the trace of the EM 3-cocycle (ω^2, c^2) can be obtained from b . It follows from (i) that ω^2 is a coboundary. $\square$

Lemma 4.5. *Let G be a finite abelian group and (ω, c) an EM 3-cocycle of G . Then ω is a coboundary if, and only if, $\bar{\nu}(V) > 0$ for all simple $H(G, \omega)$ -module V . In this case, $H(G, \omega)$ is gauge equivalent to the Hopf algebra $\mathbb{C}[G]^*$.*

Proof. In view of Example 2.5, if ω is a coboundary, then $H(G, \omega)$ can be twisted to the bialgebra $\mathbb{C}[G]^*$ by a gauge transformation. It follows from Corollary 3.5 that $\bar{\nu}(V) > 0$ for all simple $H(G, \omega)$ -modules V . Conversely, we assume the positivity of total indicators. Let t be the trace of the EM 3-cocycle (ω, c) . By Lemma 4.4(i), it is equivalent to show that $t \in K(G, \mathbb{C}^\times)$. Since G is a direct sum of its cyclic subgroups, by [23, Lem. 6.2(i)], $t \in K(G, \mathbb{C}^\times)$ if, and only if, $t_C \in K(C, \mathbb{C}^\times)$ for each cyclic summand C of G , where t_C denotes the restriction of t on C . Note that the restriction (ω_C, c_C) of (ω, c) on C is an EM 3-cocycle of C and its trace is equal to t_C . Therefore, by Lemma 4.4(i), it is enough to prove that ω_C is a coboundary for each cyclic subgroup C of G .

By Lemma 4.4(ii), ω^2 is a coboundary. We may simply assume $\omega^2 = 1$ as $\bar{\nu}(V)$ are preserved by gauge equivalence of semisimple quasi-Hopf algebras. A complete set of non-isomorphic simple $H(G, \omega)$ -modules is given by $\text{Irr}(H(G, \omega)) = \{V_x | x \in G\}$. By [25, Prop. 7.1],

$$(4.2) \quad \nu_n(V_x) = \delta_{x^n, 1} \prod_{r=1}^{n-1} \omega(x, x^r, x).$$

The equation implies that $\nu_n(V_x) = 0$ if $\ell \nmid n$ where $\ell = \text{ord}(x)$. Since $\omega^2 = 1$, $\omega(x, x^r, x) = \pm 1$ for all $r \in \mathbb{Z}$. By (4.2), $\nu_\ell(V_x) = \pm 1$ and

$$\nu_{k\ell}(V_x) = \nu_\ell(V_x)^k.$$

for all positive integer k . We claim that $\nu_\ell(V_x) = 1$. Suppose not. Then $\nu_\ell(V_x) = -1$. Let $N = \text{FSexp}(H(G, \omega))$. Then $\nu_N(V_x) = \dim V_x = 1$. Therefore, N/ℓ is even and so $\bar{\nu}(V_x) = 0$ which contradicts the assumption of the positivity of total indicators.

We now have

$$1 = \nu_\ell(V_x) = \prod_{r=1}^{\ell-1} \omega(x, x^r, x).$$

Note that right hand side is also the ℓ -th indicator of V_x considered as an $H(C, \omega_C)$ -module where C is the cyclic subgroup generated by x . Since $\nu_\ell(V_x)$ is invariant under gauge transformations, the product depends only on the cohomology class of ω_C in $H^3(C, \mathbb{C}^\times)$, which is a cyclic group of order ℓ . A generating 3-cocycle ϕ , as described in [20], is defined by

$$\phi(x^i, x^j, x^k) = q^{\bar{i}(\bar{j} + \bar{k} - \overline{j+k})/\ell}$$

where $\bar{i}$ denotes the least non-negative residue of i modulo ℓ , and q a primitive ℓ -th root of unity. Thus, ω_C is cohomologous to ϕ^a for some non-negative integer $0 \leq a \leq \ell - 1$, and

$$1 = \prod_{r=1}^{\ell-1} q^{a(r+1-\overline{r+1})/\ell} = q^a.$$

This implies $a = 0$ and so ω_C is a coboundary. This completes the proof of the lemma. $\square$

Now we can proof Theorem 4.1

Proof Theorem 4.1. By [23, Cor. 3.6], $D^\omega(G)$ is spanned by the set Γ^ω of group-like elements when $D^\omega(G)$ is commutative. In particular, Γ^ω is a finite abelian group which fits into an exact sequence of abelian groups

$$1 \rightarrow \hat{G} \rightarrow \Gamma^\omega \rightarrow G \rightarrow 1$$

determined by G and ω . Moreover, $D^\omega(G)$ is isomorphic to $H(\Gamma^\omega, \omega')$ as quasi-bialgebras, where $\omega' \in Z^3(\Gamma^\omega, \mathbb{C}^\times)$ is the inflation of ω^{-1} along the above map $\Gamma^\omega \rightarrow G$ [23, Sect. 9]. The braiding of $\text{Rep}(D^\omega(G))$ determines an EM 3-cocycle (ω', c) of Γ^ω . By Lemma 4.5, the existence of a vanishing total indicator if, and only if, ω' is a non-trivial 3-cocycle of Γ^ω . This is equivalent to that $H(\Gamma^\omega, \omega')$, or equivalently $D^\omega(G)$, is a genuine quasi-Hopf algebra. $\square$

5. TAMBARA-YAMAGAMI CATEGORIES

In this section, we study the positivity of total indicators for *integral* Tambara-Yamagami categories associated with elementary p -groups [31]. By [8], these fusion categories are monoidally equivalent to the categories of representations of semisimple quasi-Hopf algebras. In [32], a necessary and sufficient condition is obtained for an integral Tambara-Yamagami category admitting a fibration, i.e. it is monoidally equivalent to the category of representations of a semisimple Hopf algebra. By Corollary 3.5, positivity of total indicators is a necessary condition for the existence of a fibration, but not sufficient in general. We will demonstrate the sufficiency of positivity of total indicators for these integral Tambara-Yamagami categories in this section.

Let A be a finite abelian group. Tambara and Yamagami [31] classified fusion categories with a complete set of simple objects $A \sqcup \{m\}$ satisfying the fusion rules

$$(5.1) \quad a \otimes b \cong ab, \quad a \otimes m \cong m \cong m \otimes a, \quad m \otimes m \cong \bigoplus_{x \in A} x \quad (a, b \in A).$$

They also showed that such categories are parametrized by pairs (χ, τ) where χ is a non-degenerate symmetric bicharacter of A , and τ is a square root of $|A|^{-1}$. The corresponding fusion category, denoted by $\mathcal{TV}(A, \chi, \tau)$, can be described more precisely in the following definition.

Definition 5.1. $\mathcal{TV}(A, \chi, \tau)$ is a skeletal fusion category over $\mathbb{C}$ with the set of simple objects $S := A \sqcup \{m\}$. Hom-sets between elements of S are given by

$$\text{Hom}(s, s') = \begin{cases} \mathbb{C} & \text{if } s = s', \\ 0 & \text{otherwise} \end{cases}$$

with $\text{id}_s = 1 \in \mathbb{C}$. The compositions of morphisms are obvious one. Tensor products of elements of S are given by (5.1) (but with $\cong$ replaced by $=$). The unit object $\mathbf{1}$ is strict and equal to the identity $e \in A$. The associativity constraint Φ is determined by

$$\begin{aligned} \Phi_{a,m,b} &= \chi(a, b) \text{id}_m : m \rightarrow m, \\ \Phi_{m,a,m} &= (\chi(a, x) \delta_{x,y} \text{id}_x)_{x,y} : \bigoplus_{x \in A} x \rightarrow \bigoplus_{y \in A} y, \\ \Phi_{m,m,m} &= (\tau \chi(x, y)^{-1} \text{id}_m)_{x,y} : \bigoplus_{x \in A} m \rightarrow \bigoplus_{y \in A} m, \end{aligned}$$

where $a, b \in A$, and the other $\Phi_{s,t,u}$ ($s, t, u \in S$) are identity morphisms. Duality of $\mathcal{TV}(A, \chi, \tau)$ is described as follows: For $a \in A$, $a^* := a^{-1}$ with ev_a, db_a being identity id_e . For the object m , $m^* := m$ with morphisms $\text{ev}_m = \tau^{-1} \pi : m \otimes m \rightarrow \mathbf{1}$ and $\text{db}_m = \iota : \mathbf{1} \rightarrow m \otimes m$ where $\pi : m \otimes m \rightarrow \mathbf{1}$ and $\iota : \mathbf{1} \rightarrow m \otimes m$ are respectively the canonical projection and embedding satisfying $\pi \iota = \text{id}_{\mathbf{1}}$

Since the Frobenius-Perron dimension of $\mathcal{TV}(A, \chi, \tau)$ is an integer, $\mathcal{TV}(A, \chi, \tau)$ is pseudo-unitary by [8]. In particular, there exists a unique spherical pivotal structure j for which the pivotal dimension $d(V)$ of an object V is equal to its Frobenius-Perron dimension, i.e.

$$d(a) = 1 \quad \text{for } a \in A, \quad \text{and} \quad d(m) = \sqrt{|A|}.$$

One can verify directly that $j_a = \text{id}_a$ for $a \in A$, and $j_m = \text{sgn}(\tau) \text{id}_m$, where $\text{sgn}(\tau)$ means the sign of the real number τ . We will always assume $\mathcal{TV}(A, \chi, \tau)$ is a spherical fusion category relative to this canonical pivotal structure j .

By [31], two Tambara-Yamagami categories $\mathcal{TV}(A, \chi, \tau)$ and $\mathcal{TV}(A', \chi', \tau')$ are monoidally equivalent if, and only if, $\tau = \tau'$ and there exists a group isomorphism $\sigma : A \rightarrow A'$ satisfying $\chi(a, b) = \chi'(\sigma(a), \sigma(b))$ for all $a, b \in A$ (i.e. (A, χ) and (A', χ') are *isometric*).

We now turn to the case of elementary p -groups V of order p^r , where p is a prime. Then V is an r -dimensional vector space over the finite field $\mathbb{F}_p$ of order p . A non-degenerate symmetric bilinear form $B : V \times V \rightarrow \mathbb{F}_p$ on V determines a non-degenerate symmetric bicharacter χ_B defined by $\chi_B(a, b) = \exp(2\pi i B(a, b))$; moreover, the assignment $B \rightarrow \chi_B$ is a one-to-one correspondence. Two bilinear forms B and B' respectively on the $\mathbb{F}_p$ -spaces V and V' are said to be *isometric* if there exists an isomorphism $\sigma : V \rightarrow V'$ of $\mathbb{F}_p$ -spaces such that $B'(\sigma(a), \sigma(b)) = B(a, b)$ for all $a, b \in V_1$. It is easy to see that two bilinear forms B, B' are isometric if, and only if, $\chi_B, \chi_{B'}$ are isometric bicharacters. Moreover, any bilinear form B on a $\mathbb{F}_p$ -space is uniquely determined by its *Gram matrix* $[B(v_i, v_j)]_{i,j}$ relative to a basis $\{v_i\}_i$ of V . In particular, we will denote by B_0 the bilinear form on $\mathbb{F}_p^r$ whose Gram matrix relative to the standard basis is the identity

Remark 5.2. The Tambara-Yamagami category $\mathcal{TV}(A, \chi, \tau)$ is integral if, and only if, $d(m) = \sqrt{|A|}$ is an integer, or equivalently $|A|$ is a square. In this case, by [8], the fusion category is monoidally equivalent the representation category of

a semisimple quasi-Hopf algebra over $\mathbb{C}$. If V is an elementary p -group of order p^r , then $\mathcal{TY}(V, \chi, \tau)$ is integral if, and only if, r is even.

5.1. Characteristic two. We will show in this subsection that a Tambara-Yamagami category associated with an elementary 2-group of square order admits a fibration if, and only if, all its total indicators are positive.

Recall that all the non-degenerate alternating bilinear forms on $V = \mathbb{F}_2^r$ with r even are isometric. Any non-degenerate symmetric bilinear form on V , which is not alternating, is isometric to B_0 . In particular, there are exactly two isometric classes of non-degenerate symmetric bilinear on $\mathbb{F}_2^r$ when r is even. For odd r , every non-degenerate symmetric bilinear form is isometric to B_0 (see, for example, [1]).

Using this classification of symmetric bilinear forms, the indicators of the object m in these Tambara-Yamagami categories have been obtained by Shimizu [30, Thm. 6.3]: For any non-degenerate symmetric bilinear form B on $\mathbb{F}_2^r$, the n -th indicator $\nu_n(m)$ of the simple object m in $\mathcal{TY}(\mathbb{F}_2^r, \chi_B, \tau)$ is zero if n odd. Moreover,

(i) if B is not alternating, then

$$(5.2) \quad \nu_{2k}(m) = \text{sgn}(\tau)^k \left(\frac{1+i}{\sqrt{2}} \right)^{rk} \left(\frac{1+i^{-k}}{\sqrt{2}} \right)^r.$$

(ii) If r is even and B is alternating, then

$$(5.3) \quad \nu_{2k}(m) = \begin{cases} \text{sgn}(\tau) & \text{if } k \text{ is odd,} \\ 2^{r/2} & \text{if } k \text{ is even.} \end{cases}$$

We can now compute the total indicators for the integral Tambara-Yamagami categories associated with an elementary 2-group of square order.

Proposition 5.3. *Let $V = \mathbb{F}_2^{2\ell}$, B a non-degenerate symmetric bilinear form on V , τ a square root of $|V|^{-1}$, and $\mathcal{C} = \mathcal{TY}(V, \chi_B, \tau)$.*

(i) *If B is not alternating, then $\text{FSexp}(\mathcal{C}) = 8$ and $\bar{\nu}(m) = 2\text{sgn}(\tau) + 2^\ell$.*

(ii) *If B is alternating, then $\text{FSexp}(\mathcal{C}) = 4$ and $\bar{\nu}(m) = \text{sgn}(\tau) + 2^\ell$.*

In particular, $\bar{\nu}(m) = 0$ if, and only if, B is not alternating, $\ell = 1$ and $\text{sgn}(\tau) = -1$.

Proof. Note that for all $a \in V$, $\nu_n(a) = 1 = d(a)$ if n is even, and 0 otherwise.

(i) If B is not alternating, then, by (5.2), we have

$$\nu_2(m) = \text{sgn}(\tau), \quad \nu_4(m) = 0, \quad \nu_6(m) = \text{sgn}(\tau), \quad \nu_8(m) = 2^\ell = d(m).$$

Therefore, $\text{FSexp}(\mathcal{C}) = 8$ and $\bar{\nu}(m) = \sum_{k=1}^4 \nu_{2k}(m) = 2\text{sgn}(\tau) + 2^\ell$.

(i) If B is alternating, then, by (5.3), we have

$$\nu_2(m) = \text{sgn}(\tau) \quad \text{and} \quad \nu_4(m) = 2^\ell = d(m).$$

Therefore, $\text{FSexp}(\mathcal{C}) = 4$ and $\bar{\nu}(m) = \sum_{k=1}^2 \nu_{2k}(m) = \text{sgn}(\tau) + 2^\ell$.

The last statement follows directly from statement (i) and (ii). $\square$

Corollary 5.4. *Let $\mathcal{C}$ be a Tambara-Yamagami category associated with an elementary 2-group V of order $2^{2\ell}$. Then $\mathcal{C}$ is monoidally equivalent to $\text{Rep}(H)$ for some semisimple Hopf algebra H if, and only if, all its total indicators are positive.*

Proof. By [32, Prop 5.5], $\mathcal{C}$ has a fibration if, and only if, $\mathcal{C}$ is not monoidally equivalent to $\mathcal{TY}(\mathbb{F}_2^2, B_0, -2)$. It follows from Proposition 5.3 that the simple object m in $\mathcal{C}$ satisfies $\bar{\nu}(m) > 0$ if, and only if, $\mathcal{C}$ is not monoidally equivalent to $\mathcal{TY}(\mathbb{F}_2^2, B_0, -2)$. This proves the corollary. $\square$

5.2. Odd characteristic. In contrast to the characteristic two case, positivity of total indicators may not be sufficient for the existence of fibration of an integral Tambara-Yamagami category associated with an elementary p -group for odd p . We will demonstrate this fact by computing the total indicators. By [29, Ch. 4], there are exactly two isometric classes of non-degenerate symmetric bilinear form on an r -dimensional $\mathbb{F}_p$ -space. They are represented by B_0 and B_1 whose Gram matrix relative to the standard basis is given by

$$\left[\begin{array}{c|c} I_{r-1} & 0 \\ \hline 0 & u \end{array} \right]$$

where u can be any fixed quadratic nonresidue in $\mathbb{F}_p$, and I_{r-1} denotes the identity matrix of rank $r-1$. In particular, a non-degenerate symmetric bilinear form B on $\mathbb{F}_p^r$ is determined by its *discriminant* $\det(B)$ in $\mathbb{F}_p^\times/(\mathbb{F}_p^\times)^2$, where

$$\det B = \det ([B(e_i, e_j)]_{ij}) \in \mathbb{F}_p^\times.$$

Therefore, the isometric class of a bilinear form B is uniquely determined by the Legendre symbol $\left(\frac{\det B}{p}\right)$ which is 1 if $\det B$ is a quadratic residue, and -1 otherwise. We can now compute the total indicators using the following formula obtained by Shimizu [30, Thm. 6.1]: The n -th indicator of m in $\mathcal{TY}(\mathbb{F}_p^r, \chi_B, \tau)$ for any non-degenerate symmetric bilinear form B on $\mathbb{F}_p^r$ is given by

$$(5.4) \quad \nu_n(m) = \begin{cases} 0 & \text{if } n \text{ is odd,} \\ \text{sgn}(\tau)^k \varepsilon_p^{r(k+1)} \left(\frac{-k}{p}\right)^r \left(\frac{-2}{p}\right)^{r(k+1)} \left(\frac{\det B}{p}\right)^{k+1} & \text{if } n = 2k \text{ and } p \nmid k, \\ \text{sgn}(\tau)^k \varepsilon_p^{rk} \left(\frac{-2}{p}\right)^{rk} \left(\frac{\det B}{p}\right)^k \sqrt{p^r} & \text{if } n = 2k \text{ and } p \mid k, \end{cases}$$

where $\varepsilon_p = \sqrt{\left(\frac{-1}{p}\right)}$.

Proposition 5.5. *Let $V = \mathbb{F}_p^{2\ell}$, χ_B the bicharacter associated with a non-degenerate bilinear form B on V , τ a square root of $|V|^{-1}$, and $\mathcal{C} = \mathcal{TY}(V, \chi_B, \tau)$.*

(i) *If $\left(\frac{-1}{p}\right)^\ell \left(\frac{\det B}{p}\right) \text{sgn}(\tau) = 1$, then $\text{FSexp}(\mathcal{C}) = 2p$ and*

$$\bar{\nu}(m) = p^\ell + (p-1) \text{sgn}(\tau).$$

(ii) *If $\left(\frac{-1}{p}\right)^\ell \left(\frac{\det B}{p}\right) \text{sgn}(\tau) = -1$, then $\text{FSexp}(\mathcal{C}) = 4p$ and $\bar{\nu}(m) = 0$.*

Proof. By [30, Thm. 3.2], $\nu_n(a) = \delta_{na,0}$ for $a \in V$. Therefore, $p \mid \text{FSexp}(\mathcal{C})$. To determine $\text{FSexp}(\mathcal{C})$, it is enough to consider the values $\nu_n(m)$ with $n = 2k$ and $p \mid k$ by virtue of (5.4). Note that

$$\varepsilon_p^{2\ell} \left(\frac{\det B}{p}\right) = \left(\frac{-1}{p}\right)^\ell \left(\frac{\det B}{p}\right) = \epsilon \text{sgn}(\tau)$$

for some $\epsilon = \pm 1$. Therefore, (5.4) becomes

$$(5.5) \quad \nu_n(m) = \begin{cases} 0 & \text{if } n \text{ is odd,} \\ \text{sgn}(\tau) \epsilon^{k+1} & \text{if } n = 2k \text{ and } p \nmid k, \\ \epsilon^k p^\ell & \text{if } n = 2k \text{ and } p \mid k, \end{cases}$$

If $\epsilon = 1$, then $\nu_{2p}(m) = p^\ell = d(m)$ and so $\text{FSexp}(\mathcal{C}) = 2p$. Moreover,

$$\bar{\nu}(m) = p^\ell + \sum_{k=1}^{p-1} \nu_{2k}(m) = p^\ell + \text{sgn}(\tau)(p-1).$$

If $\epsilon = -1$, then $\nu_{2p}(m) = -p^\ell$ and $\nu_{4p}(m) = p^\ell$. Thus, $\text{FSexp}(\mathcal{C}) = 4p$ and

$$\bar{\nu}(m) = \sum_{k=1}^{2p} \nu_{2k}(m) = 0. \quad \square$$

Corollary 5.6. *Let B be a non-degenerate symmetric bilinear form on $V = \mathbb{F}_p^{2\ell}$ and $\tau = \pm p^{-\ell}$. Then $\mathcal{C} = \mathcal{TY}(V, \chi_B, \tau)$ admits a fibration if, and only if, $\bar{\nu}(s) \geq d(s)$ for all simple object $s \in \mathcal{C}$.*

Proof. By the preceding proposition, for $s \in V$, $\bar{\nu}(s) \geq 2 > d(s)$. In particular, the inequality holds for all $s \in V$ automatically. Therefore, we only need to consider the simple object m .

By [32, Prop. 4.1], $\mathcal{C}$ admits a fibration if, and only if, $\tau = p^{-\ell}$ and B is hyperbolic, i.e. the Gram matrix of B relative to some basis of $\mathbb{F}_p^{2\ell}$ is of the form

$$\left[\begin{array}{c|c} 0 & I_\ell \\ \hline I_\ell & 0 \end{array} \right], \text{ or equivalently, } \left(\frac{\det B}{p} \right) = \left(\frac{-1}{p} \right)^\ell.$$

If $\mathcal{C}$ admits a fibration, then $\left(\frac{\det B}{p} \right) \left(\frac{-1}{p} \right)^\ell = 1 = \text{sgn}(\tau)$ and so $\bar{\nu}(m) = p^\ell + p - 1 > p^\ell = d(m)$ by Proposition 5.5. Conversely, if $\bar{\nu}(m) \geq p^\ell$, then, by Proposition 5.5, $\text{sgn}(\tau) = 1$ and $\left(\frac{\det B}{p} \right) \left(\frac{-1}{p} \right)^\ell = 1$, or equivalently $\left(\frac{\det B}{p} \right) = \left(\frac{-1}{p} \right)^\ell$. Therefore, $\mathcal{C}$ admits a fibration by the preceding paragraph. $\square$

Remark 5.7. The corollary implies that there exists a genuine semisimple quasi-Hopf algebra of which the total indicators of its representations are all positive. Let B be the bilinear form on $\mathbb{F}_p^{2\ell}$ whose Gram matrix relative to the standard basis is

$$\left[\begin{array}{c|c} I_{2\ell-1} & 0 \\ \hline 0 & (-1)^{\ell+1} \end{array} \right]. \text{ Then } \left(\frac{\det B}{p} \right) \left(\frac{-1}{p} \right)^\ell = -1. \text{ If we take } \text{sgn}(\tau) = -1 \text{ or } \tau = -p^{-\ell},$$

then, by Corollary 5.6 and Proposition 5.5, $\mathcal{TY}(\mathbb{F}_p^{2\ell}, \chi_B, \tau)$ is monoidally equivalent to $\text{Rep}(H)$ of a *genuine* quasi-Hopf algebra H with positive total indicators. Therefore, in general, the existence of vanishing total indicators is not a necessary condition for a semisimple quasi-Hopf algebra being genuine.

REFERENCES

- [1] A. A. Albert, Symmetric and alternate matrices in an arbitrary field. I, *Trans. Amer. Math. Soc.* 43 (1938), no. 3, 386–436. MR 1501952
- [2] B. Bakalov and A. Kirillov Jr., *Lectures on tensor categories and modular functors*. University Lecture Series, 21. American Mathematical Society, Providence, RI, 2001.
- [3] P. Bantay, The Frobenius-Schur indicator in conformal field theory. *Phys. Lett. B*, 394(1-2):87–88, 1997.
- [4] S. Burciu, On some representations of the Drinfeld double. *J. Algebra* 296 (2006), no. 2, 480–504.
- [5] R. Dijkgraaf, V. Pasquier and P. Roche, Quasi Hopf algebras, group cohomology and orbifold models. *Nuclear Phys. B Proc. Suppl.* 18B (1990), 60–72.
- [6] P. Etingof and S. Gelaki, On the exponent of finite-dimensional Hopf algebras. *Math. Res. Lett.* 6 (1999), no. 2, 131–140.
- [7] P. Etingof and S. Gelaki, On families of triangular Hopf algebras. *Int. Math. Res. Not.* (2002), no. 14, 757–768.

- [8] P. Etingof, D. Nikshych and V. Ostrik, On fusion categories. *Ann. of Math.* (2) 162 (2005), no. 2, 581-642.
- [9] C. Goff, G. Mason and S.-H. Ng, On the gauge equivalence of twisted quantum doubles of elementary abelian and extra-special 2-groups. *J. Algebra* 312 (2007), no. 2, 849-875.
- [10] A. Joyal and R. Street, Braided monoidal categories, Macquarie Math. Reports No. 850067, Dec 1985; Revised No. 860081, Nov. 1986.
- [11] Y. Kashina, On the order of the antipode of Hopf algebras in ${}^H_H\text{YD}$. *Comm. Algebra*, 27 (1999), no. 3, 1261-1273.
- [12] Y. Kashina, A generalized power map for Hopf algebras. In *Hopf algebras and quantum groups (Brussels, 1998)*, volume 209 of *Lecture Notes in Pure and Appl. Math.*, pages 159-175. Dekker, New York, 2000.
- [13] Y. Kashina, Y. Sommerhäuser and Y. Zhu, On higher Frobenius-Schur indicators. *Mem. Amer. Math. Soc.* 181 (2006), no. 855, viii+65 pp.
- [14] C. Kassel, *Quantum groups*. Springer-Verlag, New York, 1995.
- [15] R. G. Larson and D. E. Radford, Finite-dimensional cosemisimple Hopf algebras in characteristic 0 are semisimple. *J. Algebra*, 117 (1988), no. 2, 267-289.
- [16] V. Linchenko and S. Montgomery, A Frobenius-Schur theorem for Hopf algebras. Special issue dedicated to Klaus Roggenkamp on the occasion of his 60th birthday, *Algebr. Represent. Theory* 3 (2000), no. 4, 347-355.
- [17] S. MacLane, Cohomology theory of Abelian groups, *Proceedings of the International Congress of Mathematicians*, Cambridge, Mass., 1950, vol. 2, Amer. Math. Soc., Providence, R. I., 1952, pp. 8-14.
- [18] A. Masuoka, Semisimple Hopf algebras of dimension 6, 8. *Israel J. Math.* 92 (1995), no. 1-3, 361-373.
- [19] S. Montgomery, *Hopf algebras and their actions on rings*, volume 82 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1993.
- [20] G. Moore and N. Seiberg, Classical and quantum conformal field theory, *Comm. Math. Phys.* 123 (1989), no. 2, 177-254.
- [21] M. Müger, Tensor categories: a selective guided tour. *Rev. Un. Mat. Argentina* 51 (2010), no. 1, 95-163.
- [22] M. Müger, From subfactors to categories and topology. II. The quantum double of tensor categories and subfactors, *J. Pure Appl. Algebra* 180 (2003), no. 1-2, 159-219.
- [23] G. Mason and S.-H. Ng, Group cohomology and gauge equivalence of some twisted quantum doubles. *Trans. Amer. Math. Soc.* 353 (2001), no. 9, 3465-3509.
- [24] G. Mason and S.-H. Ng, Central invariants and Frobenius-Schur indicators for semisimple quasi-Hopf algebras. *Adv. Math.*, 190 (2005), no. 1, 161-195.
- [25] S.-H. Ng and P. Schauenburg, Central invariants and higher indicators for semisimple quasi-Hopf algebras. *Trans. Amer. Math. Soc.* 360 (2008), no. 4, 1839-1860.
- [26] S.-H. Ng and P. Schauenburg, Frobenius-Schur indicators and exponents of spherical categories. *Adv. Math.*, 211 (2007), no. 1, 34-71.
- [27] S.-H. Ng and P. Schauenburg, Higher Frobenius-Schur indicators for pivotal categories. *Contemporary Math.*, 441, 2007, 63-90.
- [28] S.-H. Ng and P. Schauenburg, Congruence subgroups and generalized Frobenius-Schur indicators. *Comm. Math. Phys.* 300 (2010), no. 1, 1-46.
- [29] J.-P. Serre, *A course in arithmetic*, Springer-Verlag, New York, 1973, Translated from the French, Graduate Texts in Mathematics, No. 7.
- [30] K. Shimizu, Frobenius-Schur indicators in Tambara-Yamagami categories. *J. Algebra* 332 (2011), 543-564.
- [31] D. Tambara and S. Yamagami, Tensor categories with fusion rules of self-duality for finite abelian groups. *J. Algebra* 209 (1998), no. 2, 692-707.
- [32] D. Tambara, Representations of tensor categories with fusion rules of self-duality for abelian groups. *Israel J. Math.* 118 (2000), 29-60.
- [33] V. G. Turaev, *Quantum invariants of knots and 3-manifolds*. volume 18 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1994.
- [34] C. Vafa, Toward classification of conformal theories. *Phys. Lett. B* 206 (1988), no. 3, 421-426.

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, CHINA

E-mail address: `gxliu@nju.edu.cn`

DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, AMES, IA 50011, USA

E-mail address: `rng@iastate.edu`

Current address: Department of Mathematics, Cornell University, Ithaca, NY 14850, USA